

Spectroscopic LSJ notation for atomic levels obtained from relativistic calculations

G. Gaigalas^{a,*}, T. Zalandauskas^b and S. Fritzsche^a

^a Fachbereich Physik, Universität Kassel,
Heinrich-Plett-Str. 40, D-34132 Kassel, Germany.

^b Vilnius University Research Institute of Theoretical Physics and Astronomy,
A. Goštauto 12, Vilnius 2600, Lithuania.

February 2, 2008

Abstract

Today, relativistic calculations are known to provide a very successful means in the study of open-shell atoms and ions. But although accurate atomic data are obtained from these computations, they are traditionally carried out in jj -coupling and, hence, do often not allow for a simple LSJ classification of the atomic levels as needed by experiment. In fact, this lack of providing a proper spectroscopic notation from relativistic structure calculations has recently hampered not only the spectroscopy of medium and heavy elements, but also the interpretation and analysis of inner-shell processes, for which the occurrence of additional vacancies usually leads to a very detailed fine structure.

Therefore, in order to facilitate the classification of atomic levels from such computations, here we present a program (within the RATIP environment) which help transform the atomic wave functions from jj -coupled multiconfiguration Dirac-Fock computations into a LS -coupled representation. Beside of a proper LSJ assignment to the atomic levels, the program also supports the full transformation of the wave functions if required for (nonrelativistic) computations.

*Email: gaigalas@mserv.itpa.lt

PROGRAM SUMMARY

Title of program: LSJ.

Catalogue number: ADTL

Program summary URL: <http://cpc.cs.qub.ac.uk/summaries/ADTL>

Program obtainable from: CPC Program Library, Queen's University of Belfast, N. Ireland.

Licensing provisions: None.

Computer for which the new version has been tested: IBM RS 6000, PC Pentium III.

Installations: University of Kassel (Germany).

Operating systems: IBM AIX 4.1.2+, Linux 7.1.+

Program language used in the new version: ANSI standard Fortran 90/95.

Memory required to execute with typical data: Memory requirements depend on the shell structure and the size of the wave function expansion which is used to represent the atomic levels.

No. of bits in a word: All real variables are parametrized by a `selected kind parameter` and, thus, can easily be adapted to any required precision as supported by the compiler. Presently, the `kind` parameter is set to double precision (two 32-bit words) in the module `rabs_constant`.

CPU time required to execute test data: 5 seconds on a 1 GHz Pentium III processor.

Distribution format: compressed tar file.

Keywords: atomic, *LSJ* spectroscopic notation, *LS* – *jj* transformation, multiconfiguration Dirac–Fock, recoupling of angular momenta, relativistic.

Nature of the physical problem:

The spectroscopic *LSJ* notation is determined for atomic levels which were calculated previously in the framework of the *jj*–coupled multiconfiguration Dirac–Fock (MCDF) model. This notation is based on a complete *jj* – *LS* transformation of the leading *jj*–coupled configuration state functions (CSF) in the wave function representation of the selected levels.

Restrictions onto the complexity of the problem:

The *jj* → *LS* transformation of the (*jj*–coupled) CSF is supported for all shell structures including open *s*–, *p*–, *d*–, and *f*–shells. For shells with $l > 3$ (i.e. beyond the *f*–subshells), however, a proper transformation of the antisymmetrized subshell states can be carried out only for the case of one or two *equivalent* electrons. This restriction also applies for the transformation of the $g_{7/2}$ and $g_{9/2}$ subshell states which are otherwise supported by the RATIP package [S. Fritzsche, J. Elec. Spec. Rel. Phen. **114–116** (2001) 1155.]. The *jj* ↔ *LS* transformation matrices, which are applied internally by the program, are consistent with the definition of the (reduced) coefficients of fractional parentage [G. Gaigalas, Z. Rudzikas and

C. Froese Fischer, At. Data Nucl. Data Tables, **70** (1998) 1; G. Gaigalas, S. Fritzsche and Z. Rudzikas, At. Data Nucl. Data Tables, **76** (2000) 235.] as published previously.

Unusual features of the program:

The LSJ program is designed as a part of the RATIP package [S. Fritzsche, J. Elec. Spec. Rel. Phen. **114–116** (2001) 1155.] for the computation of (relativistic) atomic transition and ionization properties. This (new) component therefore supports the transformation of all atomic states which are generated either with RATIP or by means of the GRASP92 code [F. A. Parpia, C. F. Fischer, and I. P. Grant, Comput. Phys. Commun. **94** (1996) 249.]. Moreover, the normalization of the transformed states is tested within the *LSJ*–coupled basis.

LONG WRITE-UP

1 Introduction

In atomic spectroscopy, the standard LSJ notation of the atomic levels is frequently applied for classifying the low-lying level structures of (most) atoms and ions. The knowledge of the basic LSJ symmetries of the levels often helps the interpretation of atomic spectra without detailed computations. In fact, the LSJ spectroscopic notation is found useful also in order to explain the excitation and the decay properties of individual levels, at least qualitatively. Since, in addition, most atomic structure calculations were carried out within a LS -coupled basis in the past, the LSJ designation of the levels is taken today as the *standard notation* in the literature as well as in the most data bases.

In recent years, however, relativistic structure computations also became feasible and were frequently performed, when accurate data were required. Based on either the Dirac-Coulomb or Dirac-Coulomb-Breit Hamiltonian, for instance, these computations are traditionally carried out in jj -coupling, from which the LSJ notation cannot be derived so easily, especially if atomic configurations with (several) open shells occur. In the last few years, therefore, the lack of a proper spectroscopic notation deriving from relativistic calculations has hampered not only further analysis of open d - and f -shell elements but also the study and interpretation of the inner-shell processes where the occurrence of (additional) inner-shell holes often result in a very detailed fine-structure. In order to obtain the standard notation in such a case, an additional transformation of the atomic states (from the jj -coupled into a LS -coupled basis) is needed first before the full benefit is drawn from such a relativistic treatment. But although several tools and tabulations are available today for carrying out the transformation between different coupling schemes, there is no program available which would directly support the well-known relativistic structure codes such as GRASP92 [1] or RATIP [2, 3, 4]. The reasons for this lack are simple: (i) the rapidly increasing complexity of the transformation matrices, when several open shells are involved, and (ii) no simple access to the standard quantities from the atomic shell model.

Apart from the explicit $jj - LS$ transformation of the many-electron basis, there is a second method known from the literature [5, 6, 7] which helps to derive the LSJ classification of atomic levels from relativistic computations. This method applies the (*effective*) relativistic operators within an LS -coupled basis and, hence, can be directly implemented into the non-relativistic theory, i.e. by using the familiar Pauli spinors for the construction of the many-electron basis. So far, however, these effective operators have neither been derived in detail for all the physical cases of interest [8] nor does a full implementation exist which is competitive with the standard relativistic codes. Moreover, the use of effective operators within a Pauli spinor basis cannot account for the different *radial* behaviour of equivalent electrons with $j = l \pm 1/2$ ($l \geq 1$) as

is found to be important, in particular, for the behaviour of inner-shell electrons in medium and heavy elements. Therefore, the explicit $jj - LS$ transformation of the many-electron basis seems to us a much simpler route to derive the LSJ designation of the atomic states, making use of the well-known relativistic structure codes such as GRASP92 [1] or RATIP [2] as its input. This $jj - LS$ transformation of the basis states has recently been generalized also to allow an arbitrary number of open shells in the construction of the symmetry-adapted functions [9].

To facilitate the classification of the atomic levels and wave functions from the relativistic calculations, here we present the LSJ program as a new component of the RATIP package [2]. This component supports both the LSJ classification of the level structure of open-shell atoms and ions as well as the (complete) transformation of the corresponding wave functions. But the program may also be of value in solving the problems of a more general character, where the transformation from the jj - into LS -coupling scheme and vice versa is performed, e.g. a transformation of this type is needed when one seeks the optimal coupling scheme after calculating the energy spectra in a non-relativistic or relativistic approximation. But the problems of this type are quite independent of those investigated here. They are related only to a similar transformation being made. Therefore the authors have not aimed at creating a general program in the frame of the RATIP package that would be of use while the physical problems are formulated in the most general case, especially with certain risks originating from such a formulation of a problem when the transformation considered is based on the transformation coefficients [10]. The values of these depend on the phase convention that is chosen to obtain the spin-angular coefficients and on the definition of the coefficients of fractional parentage (CFP). Of course, the program presented in this work may serve as a basis for solving the other problems in atomic physics. But the authors have not aimed at the exhaustive description of the program on purpose, because in that case the technical way of answering those questions is not as important as is the methodological one, related to the phase convention and the definition of CFP. The program is based on the methodology that uses the Fano-Racah convention, and the CFPs are defined via the quasispin formalism [11]. The CFPs are published in [12, 13]. In all other cases this transformation leads to erroneous results. Therefore, before starting to perform any tasks related to the use of the program in other packages or trying to solve any problems of other type, we recommend that the authors of the paper should be contacted.

In the following section, we first provide a brief outline on the transformation of the atomic states in going from the jj -coupled to a LS -coupled many-electron basis. This shows how the LSJ spectroscopic notation is obtained as a symmetry of the leading term in the wave function representation within the LS -coupled basis; however, we leave out most of the details concerning the construction and recoupling of the symmetry functions in the atomic shell model since that material has been presented recently [14]. In section 3, we describe the program organization and how the code is distributed. To demonstrate the interactive use of the LSJ program, section 4 displays — by two examples — the derivation of the LSJ notation for the low-lying excited levels of beryllium as well as for the ground-state of gadolinium. Finally, the conclusions and a brief outlook are given in section 5.

2 Theoretical background

The task to derive the LSJ designation of atomic levels from relativistic calculations can be summarized rather easily without much knowledge about the atomic shell model. Having, for example, a representation of an atomic state ψ with respect to a jj -coupled symmetry basis, the (first few) leading terms in the wave function expansions need to be transformed into a $LS(J)$ -coupled basis from which, then, the proper spectroscopic notation is given by the symmetry of (again) the leading term. At the same time, the contributions from the subsequent terms in the LS -coupled basis indicate the admixture of other LS symmetries and, hence, the deviation from a pure LS or Russell-Saunders coupling scheme. Of course, the explicit $jj - LS$ transformation of the atomic state functions (ASF) requires quite detailed information about the construction and the coupling of the symmetry adapted basis functions, both in jj - as well as LS -coupling. For most practical applications, however, these details are irrelevant.

In this write-up of the LSJ program, we therefore restrict the discussion mainly to the representation of the atomic states in the various coupling schemes, but otherwise refer the reader to the literature [14, 15]. While this restriction enables us to explain the basis steps of a $jj - LS$ transformation of atomic states from standard relativistic computations, and how these steps are implemented within the program, it avoids the need of providing too many details about the definition of symmetry-adapted functions and about the theory of transformation in the atomic shell model.

2.1 Representation of atomic states

In principle, of course, any valid coupling scheme can be used in order to represent the (many-electron) wave functions in atomic structure computations. In relativistic calculations, however, a jj -coupled basis appears particularly appropriate as the symmetry of the Dirac equation for a single electron leads naturally to solutions which have a well-defined total angular momentum, $\mathbf{j} = \mathbf{l} + \mathbf{s}$, but no defined projections (onto the z -axis) of either the orbital angular momentum $\mathbf{l}$ nor the spin $\mathbf{s}$ of the electron. In the multiconfiguration Dirac-Fock (MCDF) or configuration interaction (CI) methods, these one-electron solutions of the Dirac equation (which are finally determined for some effective potential) are then used also for the set-up of the many-electron basis functions, the so-called configuration state functions (CSF). Owing to the overall symmetry of atoms with respect to a rotation or inversion of the coordinates, these CSF are constructed in order to have a well-defined total angular momentum J and total parity P . For a given set of such configuration states, $\{|\gamma_r JP\rangle\}$, $r = 1 \dots n_c$, the atomic states are obtained simply as a linear combination

$$|\Psi_\tau(J^P)\rangle = \sum_r^{n_c} a_r^{(jj)}(\tau) |\gamma_r JP\rangle \quad (1)$$

of CSF of the same symmetry, where $\tau = 1, 2, \dots$ enumerates the atomic levels (of the given symmetry) and γ_r denote the set of all further quantum numbers as required for an unique classification of the symmetry-adapted basis. Obviously, these quantum numbers and the details of their coupling play an important role in the transformation of the basis from one coupling scheme into another. For the present discussion, however, it is sufficient to say that the CSF basis in (1) spans a (finite) part of the N -electron Hilbert space and that the $a_r^{(jj)}(\tau)$

represent the (usual) Fourier coefficients for the expansion of an atomic state with respect to this basis. As seen from ansatz (1), moreover, only J and P are in fact good quantum numbers which, apart from the subshell angular momenta (see below) and the total energy of the ASF, can be used for the classification of the atomic levels.

An (almost) equivalent¹ but alternative representation of the ASF can be obtained within a LS -coupled basis

$$|\Psi_\tau(J^P)\rangle = \sum_s^{n_d} a_s^{(LS)}(\tau) |\gamma_s LS JP\rangle, \quad (2)$$

where one-electron spin-orbital functions with well-defined orbital angular momentum l (as well as with projections m_l and m_s of the orbital angular momentum and the spin onto the quantization axis) are used in the set-up of the corresponding CSF basis. Instead of a defined total angular momentum j of each single electron, however, here the orbital angular momenta and the spins of all electrons are first coupled together separately in order to obtain a total orbital angular momentum, $\mathbf{L} = \mathbf{l}_1 + \mathbf{l}_2 + \dots$, and a total spin, $\mathbf{S} = \mathbf{s}_1 + \mathbf{s}_2 + \dots$. The symmetry J of the CSF with respect to a rotation of the coordinates are obtained then from the coupling $\mathbf{J} = \mathbf{L} + \mathbf{S}$ of the total orbital and spin angular momenta. In LS -coupling, therefore, L and S are (approximately) also *good* quantum numbers which can be used for the classification of the atomic states. And, in fact, such an LS coupling of the atomic levels appears to be well realized in Nature, in particular for light atoms and ions with, say, $Z \lesssim 20$ as well as for the low-lying levels of most neutral atoms.

As pointed out, the two representations (1) and (2) of the atomic states are almost equally suitable to determine the energies and properties of an atom. To derive the LSJ designation of an atomic state, for which a representation $\{a_r^{(jj)}\}$ in jj -coupling is given, it is sufficient to expand each CSF $|\gamma_r JP\rangle$ into a LS -coupled basis and to sum up the contributions of the different CSF, using the proper weight $a_r^{(jj)}(\tau)$. For two orthonormalized sets of CSF, such a transformation

$$|\gamma_r LS JP\rangle \longleftrightarrow |\gamma_s JP\rangle \quad (3)$$

is represented by an *orthogonal* matrix. The knowledge of this matrix is therefore all that is needed in practice in order to carry out the transformation explicitly and to find the symmetry of the leading term in the LS -coupled basis. The LSJ program enables the user to evaluate and to apply this matrix for any (given) configuration basis in jj -coupling which is used for the description of open-shell atoms and ions. There are, however, a few restrictions on the set-up of these transformation matrices which are reasonable to *maintain* for practical computations. The next subsection explains these restrictions and summarizes the steps which are needed to construct the transformation matrix.

2.2 Transformation of configuration symmetry functions

For identical fermions, the symmetry-adapted CSF must represent many-electron functions which are also antisymmetric under the interchange of any pair of electron coordinates. In the

¹Since a jj -coupled basis allows, due to distinctions of *equivalent* electrons with $j = l \pm 1/2$ ($l \geq 1$), a somewhat greater flexibility in the representation of the atomic states, a *full* equivalence of the representations in jj - and LS -coupling is obtained only in the limit of a complete, infinitely large basis.

atomic (and nuclear) shell model, therefore, it is common practice to construct the CSF from antisymmetrized (sub-)shell states which describe N *equivalent* electrons in a given shell. In LS -coupling, for example, such subshell states for the shell (nl) are written as [15]

$$|nl^N \alpha \nu LS\rangle, \quad (4)$$

where n is the principal quantum number and α represents all those quantum numbers which, in addition to the total orbital angular momentum L , total spin S and seniority number ν , are needed for the unique classification of these states. In practice, an additional number α is required only for shells with orbital angular momenta $l \geq 3$, i.e. for electrons from the f -, g -, ...shells. A more detailed discussion of all the possible subshell states (for shells with $l \leq 3$) has been given recently in Refs. [13, 16]. The CSF with u shells in LS -coupling takes the form

$$\begin{aligned} & |\gamma LS JP\rangle \\ & \equiv \left| n_1 l_1^{N_1} n_2 l_2^{N_2} \dots n_u l_u^{N_u} \alpha_1 \nu_1 L_1 S_1 \dots \alpha_u \nu_u L_u S_u L_{12} S_{12} \dots L_{u-1} S_{u-1} LS J \right\rangle. \end{aligned} \quad (5)$$

The *additional* intermediate angular momenta L_{ij} , S_{ij} will arise in (5) if the subshell states of two or more open shells are coupled to each other as needed for CSF with more than a single open shell.

In jj -coupling, similarly, the subshell states of N equivalent electrons of a subshell $(n\kappa)$ are represented by [14]

$$\equiv |n\kappa^N \alpha \nu J\rangle \quad (6)$$

where κ is the relativistic (angular momentum) quantum number

$$\kappa = \pm(j + 1/2) \quad \text{for} \quad l = j \pm 1/2 \quad (7)$$

and where two further quantum numbers ν and J are found sufficient in order to classify all subshell states with $j = 1/2, 3/2, 5/2, 7/2$, and $9/2$ unambiguously. An additional number α is required only for subshell states with angular momenta $j \geq 9/2$. Again, Refs. [13, 16] provide a more thorough discussion and a complete list of all allowed states for (relativistic) subshells with $j \leq 9/2$. Of course, the transformation of the subshell states lies at the heart of any transformation which is to be carried out for the symmetry-adapted CSF and which need to be solved first. The $jj \leftrightarrow LS$ transformation of the subshell states (in both directions) was therefore the topic of a recent extension of the RACAH package [14, 17] which was utilized also for the implementation of the present program. In going from a jj - to a LS -coupled basis, however, there are two difficulties in the transformation procedures which one must keep in mind. These difficulties are related (i) to the group-theoretical differences of the subshell states in jj - and LS -coupling, respectively, as well as (ii) to the re-coupling of the antisymmetrized subshell states which is generally required in order to bring these states into a pre-defined coupling order on both sides of the transformation matrix elements. We will briefly discuss, in turn, these two issues to make the reader aware also of the limitations of the present program.

Although, in LS - and jj -coupling, the basic notations (4) and (6) of the subshell states appear to be rather similar, these states generally belong to different irreducible representations of the SO_3 rotation group because, in jj -coupling, each (nonrelativistic) nl -shell can be ‘split’

into two (relativistic) subshells with total angular momenta $j_{\pm} = l \pm 1/2$. In the expansion of a single shell state $|nl^N \alpha LS\rangle$ into a jj -coupled basis, therefore, both subshell states with j_- and j_+ , i.e. $\left| \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J} \right\rangle$ and $\left| \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J} \right\rangle$, will occur. Here, again, we make use of the

relativistic quantum number κ to simplify the notation below. Obviously, $N = \bar{N} + \bar{N}$ and $\bar{\kappa} = -(\bar{\kappa} + 1) > 0$ must apply where the notation $\bar{\kappa} > 0$ and $\bar{\kappa} < 0$ has been chosen to be consistent with the corresponding total $j_{\pm} = l \pm 1/2$ and subshell angular momenta.

Making use of this notation, the transformation between the jj - and LS -coupled subshell states takes the form

$$\left| \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J}, \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J} \right\rangle J \rangle = \sum_{\alpha LS} \left| l^{(\bar{N}+\bar{N})} \alpha LS J \right\rangle \left\langle l^{(\bar{N}+\bar{N})} \alpha LS J \left| \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J}, \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J} \right\rangle J \right\rangle \quad (8)$$

including a summation over α as well as the total subshell orbital angular momentum L and the subshell spin S . As is clear from our notation in Eq. (8), $\left| \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J}, \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J} \right\rangle J \rangle$ represents a coupled state with well-defined total angular momentum J which is built from the corresponding jj -coupled subshell states with $j_{\pm} = l \pm 1/2$ and the total subshell angular momenta $\bar{J}$ and $\bar{J}$, respectively. This expansion of the jj -coupled subshell states therefore shows that, for any number of open (sub-)shells in the construction of the CSF, the transformation of these symmetry functions can always be traced back to the transformation coefficients

$$\left\langle l^{(\bar{N}+\bar{N})} \alpha LS J \left| \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J}, \begin{smallmatrix} \bar{N} \\ \bar{\kappa} \end{smallmatrix} \bar{\nu} \bar{J} \right\rangle J \right\rangle \quad (9)$$

of the individual subshell. And, in fact, these (Fourier) coefficients are the ‘building blocks’ for any transformation of the atomic states into a LS -coupled basis. They are implemented explicitly into the LSJ component (by means of a recent revision of the RACAH package [14]) since their computation requires a recursive — and rather expansive — procedure which traces back the subshell states on both sides of expression (9) to their parents states with $N - 1$ electrons, and so on. The transformation coefficients (9) are implemented for all (sub-) shells with $l \leq 3$ and occupation numbers $N = 1, 2, \dots, 2l + 1$; for all higher (allowed) occupation numbers $N = 2l + 2, \dots, 4l + 2$, these coefficients are simply obtained by using the *electron-hole* symmetry and the tabulated coefficients for $N' = 4l + 2 - N$ [14].

A second difficulty in the jj - LS transformation of atomic states arises from the (re-) coupling of the antisymmetrized subshell states. Since, in general, CSF with several open shells occur very frequently, a pre-defined coupling order of the individual subshell states $n_1 l_1^{N_1}, n_2 l_2^{N_2}, \dots$ need first to be specified, such as

$$\left| \left(\dots \left(\left(l_1^{N_1} \alpha_1 L_1 S_1, l_2^{N_2} \alpha_2 L_2 S_2 \right) L_{12} S_{12}, l_3^{N_3} \alpha_3 L_3 S_3 \right) L_{123} S_{123} \right) \dots \right\rangle J \rangle \quad (10)$$

for an LS -coupled basis. Of course, a similar coupling order of the subshell states could be written also for a jj -coupled basis $\left| \kappa_1^{N_1} \nu_1 J_1 \right\rangle, \left| \kappa_2^{N_2} \nu_2 J_2 \right\rangle, \dots$ where any (predefined) sequence would give rise to a valid many-particle basis. But the predefined sequence of jj -coupled subshell states also decides how many (re-) coupling steps are needed to bring both sides in the overall transformation matrix $\langle \gamma_s LS JP | \gamma_r JP \rangle$ into the same coupling

order. For this reason, we request the jj -coupled CSF in the LSJ program to be given in a (so-called) *standard order*

$$\left| \left(\dots \left(\left(\left(\left(\bar{\kappa}_1^- \bar{\nu}_1^- J_1, \bar{\kappa}_1^+ \bar{\nu}_1^+ J_1 \right) J_1, \bar{\kappa}_2^- \bar{\nu}_2^- J_2 \right) J_2, \bar{\kappa}_2^+ \bar{\nu}_2^+ J_2 \right) J_2, \bar{\kappa}_3^- \bar{\nu}_3^- J_3 \right) J_3, \bar{\kappa}_3^+ \bar{\nu}_3^+ J_3 \right) J_{123} \dots \right) J \right\rangle \quad (11)$$

in which these functions have to fulfill two additional conditions:

- (i) If both subshells with common l_i , i.e. $\bar{\kappa}_i^-$ and $\bar{\kappa}_i^+$, appear in the expansion, these two subshells always occur successively in the sequence $(\bar{\kappa}_i^- \bar{\nu}_i^- J_i, \bar{\kappa}_i^+ \bar{\nu}_i^+ J_i) J_i$. Formally, we can use this sequence even for subshell states with zero occupation if we interpret $|\kappa^0 \nu = 0 \ J = 0\rangle \equiv 1$; in this case, the full Clebsch–Gordan expansion for the symbolic expression (11) remains valid due to the orthonormality properties of the Clebsch–Gordan coefficients.
- (ii) For the $jj - LS$ transformation of configuration states

$$\left\langle \left(\dots \left(\left(l_1^{N_1} \alpha_1 L_1 S_1, l_2^{N_2} \alpha_2 L_2 S_2 \right) L_{12} S_{12}, l_3^{N_3} \alpha_3 L_3 S_3 \right) L_{123} S_{123} \dots \right) J \mid \left(\dots \left(\left(\bar{\kappa}_1^- \bar{\nu}_1^- J_1, \bar{\kappa}_1^+ \bar{\nu}_1^+ J_1 \right) J_1, \bar{\kappa}_2^- \bar{\nu}_2^- J_2 \right) J_2, \bar{\kappa}_2^+ \bar{\nu}_2^+ J_2 \right) J_2, \bar{\kappa}_3^- \bar{\nu}_3^- J_3 \right) J_3, \bar{\kappa}_3^+ \bar{\nu}_3^+ J_3 \right) J_{123} \dots \right) J \right\rangle \quad (12)$$

we further assume in *standard order* that $l_1 = \bar{\kappa}_1^- = -(\bar{\kappa}_1^+ + 1)$, $l_2 = \bar{\kappa}_2^- = -(\bar{\kappa}_2^+ + 1)$, ..., i.e. that the sequence of (sub-)shell states is the *same on both sides* of the transformation matrix.

For such a standard order, explicit expression for the $jj - LS$ transformation matrix of the CSF with two open shells in LS - or up to four open shells in jj -coupling are displayed in Ref. [14]. With the present LSJ program, we now extend these expressions to allow for any number of open subshells.

Although, of course, the standard order sets a restriction onto the set-up of the jj -coupled CSF basis, it is the order which is typically applied in all relativistic structure codes such as GRASP92 [1] or RATIP [2]. But even if a *non-standard* order has been used, the LSJ component can still be utilized with only minor additional effort. Since, in fact, the coupling of the subshell states is independent of the radial wave functions, it is usually sufficient to (re-) generate the CSF basis in standard order and just to repeat the diagonalization of the Hamiltonian matrix. Of course, the representation $\{a_r^{(jj)}(\tau)\}$ of the atomic states in (1) must always be consistent with the basis itself. Moreover, if in the CSF basis the occupied shells appear in a different sequence than the usual (non-relativistic) one $1s, 2s, 2p, 3s, \dots$, the condition of the standard order is nevertheless fulfilled and the LSJ program can be applied without further modifications. In this case, however, the order of the *parent* states may become different and may finally result in a different LSJ classification, if the parent states are accounted for in the overall designation of the atomic levels.

3 Program organization and distribution

Although LSJ is provided as an independent program for the transformation of atomic levels and wave functions from a jj -coupled into a LS -coupled basis, it is designed also as a new program component of the RATIP package [2, 3, 4] and, hence, follows design principles similar

to the other components of this package. During recent years, RATIP has been found to be a powerful tool for studying the *(relativistic atomic) transition and ionization properties* of atoms and highly-charged ions within rather arbitrary configurations. Apart from the transition probabilities and lifetimes, for incorporating the interaction of atoms with the radiation field, RATIP now includes further components which help calculate Auger and photoionization properties as well as the coherence transfer in atomic cascades. As described previously, the RATIP package is built on GRASP92 [1] which is used to generate the (relativistic) wave functions within the MCDF model.

However, here we shall not recall much about RATIP itself, neither about its overall structure nor about previous applications, since this has been described previously [18]. Apart from a few further applications to incorporate atomic processes with one electron in the continuum, our recent concerns pertained to a long life-cycle of the code, making use of an object-oriented design and, in particular, the concepts of Fortran 90/95. In fact, all components of RATIP are now well prepared to deal with large wave function expansions (1) as needed for the accurate handling of open-shell systems.

3.1 Definition of derived data types

The proper choice of *derived data types* certainly belongs to the key features for a long-term maintenance and the development of a code. In RATIP, therefore, we introduced derived types at various levels of complexity, including for example a data structure `cs_function` for the representation of a single CSF (in *jj*-coupling), or the derived type `cs_basis` in order to 'store' the information about a full configuration basis. To these data structures, memory is allocated only on the basis of the actual requirements concerning the number of occupied shells and CSF in the given CSF basis. In LSJ, we now followed very similar lines and define the corresponding data structures also in *LS*-coupling. Two simple examples are

```
type, public :: cs_function_LS
  integer(kind=i1b) :: totalJ
  character(len=1)  :: parity
  integer(kind=i1b), dimension(:), pointer :: occupation
  integer(kind=i1b), dimension(:), pointer :: seniority
  integer(kind=i1b), dimension(:), pointer :: w
  integer(kind=i1b), dimension(:), pointer :: shellL
  integer(kind=i1b), dimension(:), pointer :: shellS
  integer(kind=i1b), dimension(:), pointer :: shellLX
  integer(kind=i1b), dimension(:), pointer :: shellSX
end type cs_function_LS
```

to represent a *LS*-coupled CSF in terms of the occupation and angular momentum quantum numbers of all individual subshells as well as in terms of the intermediate angular momenta $L_{12}, S_{12}, L_{123}, S_{123}, \dots$ which are stored in the two arrays `shellLX` and `shellSX`. The data relating to all the *LS*-coupled CSF are then *kept together* in the derived data type

```
type, public :: csf_basis_LS
  integer :: nocsf          ! Number of CSFs in the basis.
  integer :: nwshells       ! Number of (nonrelativistic) shells.
  integer :: nwcore         ! Number of (closed) core shells.
  integer :: number_of_electrons
  type(nl), dimension(:), pointer :: shell
  type(cs_function_LS), dimension(:), pointer :: csf
  type(parent_from_jj), dimension(:), pointer :: parent
end type csf_basis_LS
```




Figure 1: Schematic flow chart of the LSJ program.

which, within the pointer array `parent`, also includes details about the *parentage* of the corresponding configuration states functions in the original, *jj*-coupled CSF basis. Further derived structures concern the representation of the atomic states (`asf_function_LS` and `asf_basis_LS`) as well as a list of all *LS*-coupled subshell states `subshell_term_LS` to identify the allowed terms for open-shell configurations. Obviously, the efficient use of such data types facilitates the access to and the handling of selected data. In the LSJ program, all new derived types are defined in the module `rabs_lsj` (file `rabs_lsj.f90`) which also contains the transformation coefficients (9) for the *jj* ↔ *LS* transformation of the subshell states. These matrix elements are kept in the derived structure

```

type, private :: LS_jj_me
  sequence
  integer(kind=i1b)    :: w, Q, L, S, J, Nm, Qm, Jm, Qp, Jp
  integer(kind=i1b)    :: factor
  integer*8            :: nom, denom
end type LS_jj_me

```

within a *integer representation*, from which the value of the matrix element follows by computing simply: $\text{factor} \sqrt{\frac{\text{nom}}{\text{denom}}}$.

3.2 Interactive control and output of the program

Like the other components of RATIP, the LSJ program is controlled interactively. Figure 1 displays the basic steps in which the program is executed. Following the initialization of several data arrays at the beginning of the execution, the control is then taken by the procedure `lsj_control_transformation()`, which reads and interprets all input data, carries out the transformation for all selected levels, and finally prints the results to screen. This procedure also prints a *full expansion* of all *jj*-coupled CSF in the corresponding *LS*-basis in the file `input-cs1.LS`.

LSJ: Transformation of atomic states from a *jj*-coupled into a *LS*-coupled CSF basis (Fortran 95 version)
 (C) Copyright by G Gaigalas, T Zalandauskas and S Fritzsche, Kassel and Vilnius (2003).

Transform one or several ASF from a GRASP92 calculation into a *LS*-coupled CSF basis.
 The transformation starts from the given .csl and .mix files and is carried out
 for the *n* leading CSF in the *jj*-coupled basis; the new representation in the *LS* basis
 is printed similar as in a standard GRASP92 computation.

```

Enter the name of the GRASP92 configuration symmetry list file:
>
Loading configuration symmetry list file ...
  There are ... relativistic subshells;
  there are ... relativistic CSFs;
  ... load complete.
Enter the name of corresponding .mix mixing coefficient file:
>
Loading mixing coefficients file ...
  ... load complete;
Maximum number of considered ASF is:
Enter the level numbers of the ASF which are to be transformed,
e.g. 1 3 4 7 - 20 48 69 - 85 :
>
Enter the number of leading CSF to be printed:
>

```

Figure 2: Interactive dialog for LSJ at the beginning of the transformation.

Figure 2 shows the interactive dialog of LSJ which requests the user to provide the GRASP92 configuration symmetry list (.csl) file and the wave function representation (of the considered level) in a corresponding mixing coefficient (.mix) file. However, independent of the number of ASF which are represented in this .mix file, the user then needs to specify the level numbers for which the $jj \rightarrow LS$ transformation is to be carried out, as well as the number of leading *LS*-coupled CSF which are included in the printout. As discussed before, only the leading CSF is used for the *LSJ* spectroscopic designation of the level but the 'weights' of the next most important configuration states may still provide insight into the configuration mixing for the given atom or ion.

As the transformation of atomic levels appears mainly as a *geometrical task*, the definition of the configuration basis along with the corresponding mixing coefficients provide together all necessary information for a *LSJ* classification of the atomic levels. For the transformation of the corresponding wave functions, in contrast, some care must be taken also for the radial part of the wave function representation, for instance, by taking the *average* for each pair of radial orbitals with angular momentum quantum numbers $\kappa > 0$ and $-\kappa - 1$, respectively, for all occupied shells (*nl*). Using the LSJ component, the transformation of the atomic levels is carried out *promptly* for most open-shell structures; the results are therefore printed immediately to screen. Moreover, to provide a test of having obtained a *correct and complete* transformation, the sum of all 'weights' in the *LS*-coupled basis, i.e. $\sum_i |c_i^{(LS)}(\tau)|^2$ (which, of course, should be 1), is also printed.

3.3 Distribution and installation of the program

Even though the main effort in developing the LSJ component certainly concerned the two (new) modules `rabs_lsj` and `rabs_lsj_data`, we re-distribute the RATIP package as a whole, owing to the tight binding of the new components with the previously developed code. Overall, therefore, the present version of RATIP now contains 18 modules as well as the source code for the executables of the seven components ANCO, CESD99, LSJ, RCFP, RELCI, REOS99, and UTILITIES. As previously, they are all contained in the `ratip` root directory. This root also

Table I. Leading CSFs and corresponding weight coefficients ($|c_r^{(jj)}(\tau)|^2$ and $|c_s^{(LS)}(\tau)|^2$) of the low-lying levels of the $1s^2 2s 2p$ configurations in beryllium.

$ \Psi_\tau(J^P)\rangle_{jj}$		$ \Psi_\tau(J^P)\rangle_{LS}$	
weights $ c_r^{(jj)}(\tau) ^2$	CSFs	weights $ c_s^{(LS)}(\tau) ^2$	CSFs
0.66673	$ ((2s^1)_{\frac{1}{2}}, (2p^{-1})_{\frac{1}{2}}) 1\rangle$	1	$ ((2s^1)^2 S, (2p^1)^2 P) {}^3 P_1\rangle$
0.33327	$ ((2s^1)_{\frac{1}{2}}, (2p^1)_{\frac{3}{2}}) 1\rangle$		
$\sum_i c_i^{(jj)}(\tau) ^2 = 1$			$\sum_i c_i^{(LS)}(\tau) ^2 = 1$
0.66673	$ ((2s^1)_{\frac{1}{2}}, (2p^1)_{\frac{3}{2}}) 1\rangle$	1	$ ((2s^1)^2 S, (2p^1)^2 P) {}^1 P_1\rangle$
0.33327	$ ((2s^1)_{\frac{1}{2}}, (2p^{-1})_{\frac{1}{2}}) 1\rangle$		
$\sum_i c_i^{(jj)}(\tau) ^2 = 1$			$\sum_i c_i^{(LS)}(\tau) ^2 = 1$

comprises several makefiles for generating the executable as well as a number of test suites for a few components. From this root, the executables of the different components are generated by running the command `make -f make-component`, that is `make -f make-lsj` in the present case.

By utilizing a large set of different makefiles, it has been found convenient to declare several global variables for the compilation and linkage of the program. These *declarations* are now comprised within the script file `make-environment` which avoids the user having to adopt each makefile independently to the local architecture and computational environment. In fact, the script `make-environment` only contains a very few lines but must be *sourced* before, i.e. by typing `source make-environment`, to make this information available to the system. Using this set-up of the code, RATIP has been found portable rather easily to different platforms such as IBM RS/6000, SUN OS, or to the PC world. Moreover, a `Read.me` file in the `ratip` root directory contains further details to the installation.

For most components of the RATIP package, we provide in addition a test suite in a subdirectory `test-component` of the root where `component` refers to the names above. For example, the directory `test-lsj` comprises all four (input and output) files in order to run and compare the two examples below. These tests are typically designed to exhibit the capabilities of the program but may serve also as a quick test of the local installation.

4 Examples

To explain the use and execution of the LSJ component, we have selected two examples which concern (i) the well-known splitting of the beryllium terms into a singlet and triplet system and (ii) shows how the program can be used for the classification of the lanthanides with an

```

.
.
Enter the name of the GRASP92 configuration symmetry list file:
>be-csl.inp
Loading configuration symmetry list file ...
There are 4 relativistic subshells;
there are 4 relativistic CSFs;
... load complete.
Enter the name of corresponding .mix mixing coefficient file:
>be-relci.mix
Loading mixing coefficients file ...
... load complete;
Maximum number of considered ASF is: 4
Enter the level numbers of the ASF which are to be transformed,
e.g. 1 3 4 7 - 20 48 69 - 85 :
>2 4
Enter the number of leading CSF to be printed:
>2
Weights of major contributors to ASF in jj-coupling:
Level J Parity CSF contributions
2 1 - 0.66673 of 2 0.33327 of 3

Definition of leading CSF:
2) 2s ( 1) 2p-( 1)
1/2 1/2
1
Total sum of weights is: 1.0000000000000000
The new level is under the investigation.
Weights of major contributors to ASF in jj-coupling:
Level J Parity CSF contributions
4 1 - 0.66673 of 3 0.33327 of 2

Definition of leading CSF:
3) 2s ( 1) 2p ( 1)
1/2 3/2
1
Total sum of weights is: 1.0000000000000000

Weights of major contributors to ASF in LS-coupling:
Level J Parity CSF contributions
2 1 - 1.00000 of 3
Total sum of weights is: 0.9999999999999996
4 1 - 1.00000 of 2
Total sum of weights is: 0.9999999999999991

Definition of leading CSF:
2) 2s( 1) 2p( 1)
2S1 2P1 1P 1
3) 2s( 1) 2p( 1)
2S1 2P1 3P 1
LSJ complete ... .

```

Figure 3: Dialog and output for determining the LSJ notation of low-lying levels of the $1s^2 2s 2p$ configurations in beryllium.

open f -shell. In both cases, the configuration symmetry list and the corresponding mixing coefficients have been generated by means of GRASP92 [1] and the RELCI component [4] of RATIP, using simply a configuration interaction approach. The input and (expected) output of these two examples are provided also in the subdirectory `test-lsj`. In both cases, the transformation is performed very rapidly within less than a second of CPU time on a Pentium 166 MHz processor.

For beryllium, the low-lying excited spectrum of the $1s^2 2s 2p$ configuration consists of just four levels which belong to the two 1P and 3P terms and have total angular momenta $J = 1$ and $J = 0, 1, 2$, respectively. Since the Hamiltonian matrix is diagonal in the total angular momentum J , only the two levels with $J = 1$ will exhibit a configuration mixing and, in fact, this interaction is necessary to represent the two levels in terms of the $1s^2 2s^1 2p^{-1}$ and $1s^2 2s^1 2p^1$ relativistic configurations. For the two levels with $J = 0$ and $J = 2$, the assignment of the LSJ notation is trivial because there is each just one CSF with a weight coefficients equal to 1 in the expansion. But to specify also the LSJ symmetry of the remaining two $J = 1$ levels, we can run the LSJ program as shown in Figure 3, where use is made of the configuration symmetry list `be-cs1.inp` and the corresponding mixing coefficient file `be-relci.mix`. As seen from the Figure 3 and Table I where the results of the transformation are summarized, the lower one of the $J = 1$ levels (No. 2) has a 3P symmetry, and the 1P term is assigned to level No. 4. Moreover, while a strong mixing of the configuration states obviously appears in the jj -coupled basis, a (nearly) pure LSJ symmetry is found for all four levels in a LS -coupled configuration basis.

For simple shell structures, such as the low-lying excitation spectrum of beryllium and for many other light elements, the LSJ assignment can often be made just on the basis of the excitation energies (with respect to the ground state) and total angular momenta of the individual levels, if these data are compared with experiment. However, such a procedure is not feasible if two or more open shells occur in some electron configuration, or if open d - and f -shell elements are considered. In such cases, several tens or even hundreds of levels may belong to a single (nonrelativistic) configuration, making a LSJ symmetry assignment on the grounds of the calculated energies very unreliable. As a typically example for such a complex configuration, we consider the ground state of gadolinium ($Z = 64$), which has a $[\text{Xe}] 4f^7 5d 6s^2$ configuration with 3106 possible (jj -coupled) CSF. To determine the LS symmetries of the computed ground state with its known total angular momentum 2^- , we again run the LSJ component using the two files `gd-cs1.inp` and `gd-relci.mix`, respectively. Figure 4 displays the corresponding dialog and demonstrates that the ground state of gadolinium has a 9D_2 symmetry. The leading LS - and jj -coupled CSFs together with their weights (squares of mixing coefficients) are presented in Table II. The rather small mixing of other LS terms shows, moreover, that the low-lying levels of most (neutral) medium and heavy elements are still well classified in terms of their LSJ spectroscopy notation. Despite this classification, however, much larger (or even huge) wave function expansions are typically required in order to obtain accurate predictions of the level structure and properties of these systems.

5 Conclusions and outlook

The new component LSJ of the RATIP package is presented for deriving the LSJ designation of atomic levels and wave functions from relativistic calculations. Such spectroscopic notation


```

.
.
Enter the name of the GRASP92 configuration symmetry list file:
>gd-csl.inp
Loading configuration symmetry list file ...
  There are 22 relativistic subshells;
  there are 3106 relativistic CSFs;
>gd-relci.mix
... load complete.
Enter the name of corresponding .mix mixing coefficient file:

Loading mixing coefficients file ...
... load complete;
Maximum number of considered ASF is: 11
Enter the level numbers of the ASF which are to be transformed,
e.g. 1 3 4 7 - 20 48 69 - 85 :
>1
Enter the number of leading CSF to be printed:
>30
Weights of major contributors to ASF in jj-coupling:
Level J Parity      CSF contributions
  1   2   -   0.23193 of 1403  0.21537 of 457  0.12535 of 2443  0.09234 of 57  0.05331 of 987

Definition of leading CSF:
      1403) 4f-( 3)   4f ( 4)   5s ( 2)   5p-( 2)   5p ( 4)   5d-( 1)   6s ( 2)
              9/2           8
              7/2           7/2           7/2           7/2           2           2
Total sum of weights is: 0.9999999993812160

Weights of major contributors to ASF in LS-coupling:
Level J Parity      CSF contributions
  1   2   -   0.97616 of 2   0.01538 of 4   0.00483 of 6
              0.00297 of 1   0.00034 of 8   0.00005 of 100
              0.00005 of 60  0.00004 of 98  0.00004 of 58
              0.00003 of 3   0.00003 of 5   0.00002 of 102
              0.00002 of 62  0.00001 of 15  0.00001 of 28
Total sum over weight is: 0.9999999993812226

Definition of leading CSF:
  1) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      8S0 1S0 1S0 2D1 1S0 8S 8S 7D 7D 2
  2) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      8S0 1S0 1S0 2D1 1S0 8S 8S 9D 9D 2
  3) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      6P0 1S0 1S0 2D1 1S0 6P 6P 5P 5P 2
  4) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      6P0 1S0 1S0 2D1 1S0 6P 6P 7P 7P 2
  5) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      6P0 1S0 1S0 2D1 1S0 6P 6P 5D 5D 2
  6) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      6P0 1S0 1S0 2D1 1S0 6P 6P 7D 7D 2
  8) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      6P0 1S0 1S0 2D1 1S0 6P 6P 7F 7F 2
 15) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      6D0 1S0 1S0 2D1 1S0 6D 6D 7F 7F 2
 28) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      6G0 1S0 1S0 2D1 1S0 6G 6G 7D 7D 2
 58) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      4D1 1S0 1S0 2D1 1S0 4D 4D 5S 5S 2
 60) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      4D1 1S0 1S0 2D1 1S0 4D 4D 5P 5P 2
 62) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      4D1 1S0 1S0 2D1 1S0 4D 4D 5D 5D 2
 98) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      4D6 1S0 1S0 2D1 1S0 4D 4D 5S 5S 2
100) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      4D6 1S0 1S0 2D1 1S0 4D 4D 5P 5P 2
102) 4f( 7) 5s( 2) 5p( 6) 5d( 1) 6s( 2)
      4D6 1S0 1S0 2D1 1S0 4D 4D 5D 5D 2
LSJ complete ...

```

Figure 4: Dialog and output for determining the *LSJ* notation for the ground state of gadolinium.

Table II. Leading CSFs and corresponding weight coefficients ($|c_r^{(jj)}(\tau)|^2$ and $|c_s^{(LS)}(\tau)|^2$) of the ground state of gadolinium.

$ \Psi_\tau(J^P)\rangle_{jj}$		$ \Psi_\tau(J^P)\rangle_{LS}$	
weights $ c_r^{(jj)}(\tau) ^2$	CSFs	weights $ c_s^{(LS)}(\tau) ^2$	CSFs
0.23193	$ (((4f-^3\frac{9}{2}, 4f^4\ 8)\frac{7}{2}, 5d-^1\frac{3}{2})\ 2)\rangle$	0.97616	$ (4f^7\ ^8S, 5d^1\ ^2D)\ ^9D_2\rangle$
		0.01538	$ (4f^7\ ^6P, 5d^1\ ^2D)\ ^7P_2\rangle$
		0.00483	$ (4f^7\ ^6P, 5d^1\ ^2D)\ ^7D_2\rangle$
		0.00297	$ (4f^7\ ^8S, 5d^1\ ^2D)\ ^7D_2\rangle$
		0.00034	$ (4f^7\ ^6P, 5d^1\ ^2D)\ ^7F_2\rangle$
		0.00005	$ (4f^7\ ^4D^6, 5d^1\ ^2D)\ ^5P_2\rangle$
		0.00005	$ (4f^7\ ^4D^1, 5d^1\ ^2D)\ ^5P_2\rangle$
		0.00004	$ (4f^7\ ^4D^6, 5d^1\ ^2D)\ ^5S_2\rangle$
		0.00004	$ (4f^7\ ^4D^1, 5d^1\ ^2D)\ ^5S_2\rangle$
		0.00003	$ (4f^7\ ^6P, 5d^1\ ^2D)\ ^5P_2\rangle$
		0.00003	$ (4f^7\ ^6P, 5d^1\ ^2D)\ ^5D_2\rangle$
		0.00002	$ (4f^7\ ^4D^6, 5d^1\ ^2D)\ ^5D_2\rangle$
		0.00002	$ (4f^7\ ^4D^1, 5d^1\ ^2D)\ ^5D_2\rangle$
		0.00001	$ (4f^7\ ^6D, 5d^1\ ^2D)\ ^7F_2\rangle$
		0.00001	$ (4f^7\ ^6G, 5d^1\ ^2D)\ ^7D_2\rangle$
		$\sum_i c_i^{(LS)}(\tau) ^2 = 0.99998$	

is needed not only to analyze a great deal of spectroscopic data but also for the interpretation of many inner-shell phenomena such as the angular distributions of emitted Auger and photo electrons. For $2p \rightarrow 4s$ resonantly excited argon atoms, for example, it was shown recently [19] that the electron emission can be understood properly only if the decay in the (two-step) cascade is treated coherently. However, while the atomic levels in the cascade can be specified experimentally rather easily owing to the existence of observed line spectra, program tools such as LSJ are needed to understand the theoretical data and to select those levels, which have to be involved in the computation.

The design of LSJ within the framework RATIP enables the user to apply a $jj \rightarrow LS$ transformation for all atoms and shell structures which can be calculated by means of GRASP92 or any other RATIP component. This includes, in particular, open d - and f -shell elements which have attracted a lot of recent interest. For atomic fermium with $Z = 100$, for example, first measurements on the low-lying excitation spectrum have recently been carried out in Mainz and could be supported by means of extended theoretical computations [20]. Therefore, we hope that the LSJ component might be of use also in improving the available data base on the lanthanide ($Z = 57, \dots, 71$) and actinide elements ($Z = 89, \dots, 103$) in the future.

In the design and set-up of RATIP, our intention from the very beginning was to provide the user with an *open environment* for his or her studies of atomic properties. In the last few years, therefore, various requests arose for the further development of the code concerning, for example, the radiative and dielectronic recombination of highly-charged ions, electron-impact processes or even the study of multi-photon excitation and ionization processes. Very likely, however, only a few of these requests can be fulfilled by us in the forthcoming years where our attention will first be directed to the radiative capture of electrons, i.e. the reversed process to the photoionization. Therefore, in order to extend RATIP along additional lines, any support of coworkers or even the independent implementation of new components will be appreciated.

References

- [1] F. A. Parpia, C. F. Fischer and I. P. Grant, Comput. Phys. Commun. **94** (1996) 249.
- [2] S. Fritzsche, J. Elec. Spec. Rel. Phenom **114–116** (2001) 1155.
- [3] S. Fritzsche, C. F. Fischer and C. Z. Dong, Comput. Phys. Commun. **124** (2000) 342.
- [4] S. Fritzsche, C. F. Fischer and G. Gaigalas, Comput. Phys. Commun. **148** (2002) 103.
- [5] P. G. H. Sandars and J. Beck, Proc. R. Soc. London **1289** (1965) 97.
- [6] L. Armstrong, J. Math. Phys. **7** (1966) 1891.
- [7] L. Armstrong, J. Math. Phys. **9** (1968) 1083.
- [8] R. Karazija, V. Jonauskas, J. Math. Phys. **42** (2001) 5642.
- [9] G. Gaigalas, Z. Rudzikas and T. Zalandauskas, Book of Abstracts of XI HCI Conference, 1-6 September, 2002, Caen, France, p. 48.
- [10] G. Gaigalas, T. Zalandauskas and Z. Rudzikas, At. Data Nucl. Data Tables, in press (2003).
- [11] G. Gaigalas, Z. Rudzikas and C. Froese Fischer, J. Phys. **B 30** (1997) 3747.
- [12] G. Gaigalas, Z. Rudzikas and C. Froese Fischer, At. Data Nucl. Data Tables, **70** (1998) 1.
- [13] G. Gaigalas, S. Fritzsche and Z. Rudzikas, At. Data Nucl. Data Tables, **76** (2000) 235.
- [14] G. Gaigalas and S. Fritzsche, Comp. Phys. Commun. **149** (2002) 39.

- [15] Z. Rudzikas, *Theoretical Atomic Spectroscopy* (Cambridge Univ. Press, Cambridge, 1997).
- [16] G. Gaigalas, S. Fritzsche and B. Fricke, *Comp. Phys. Commun.* **135** (2001) 219.
- [17] S. Fritzsche, *Comp. Phys. Commun.* **103** (1997) 51.
- [18] S. Fritzsche, *Phys. Scr.* **T100** (2002) 37.
- [19] K. Ueda, Y. Shimizu, H. Chiba, M. Kitajima, H. Tanaka, S. Fritzsche and N. M. Kabachnik, *J. Phys.* **B 34** (2001) 107.
- [20] M. Sewtz, H. Backe, A. Dretzke, G. Kube, W. Lauth, P. Schwamb, K. Eberhardt, C. Grüning, P. Thörle, N. Trautmann, P. Kunz, J. Lassen, G. Passler, C. Z. Dong, S. Fritzsche and R. G. Haire, *Phys. Rev. Lett.*, in press (2003).